



Antimicrobial resistance targets Poland

Target achieved Progress Regress

Reduce by 27% the total consumption of antibiotics in humans



Defined daily doses (DDDs) per 1 000 inhabitants per day

At least 65% of the total consumption of antibiotics in humans belongs to the 'Access' group of antibiotics



As defined in the AWaRe classification of the WHO


*Percentage point difference from 2019.

Reduce by 10% the total incidence of bloodstream infections with meticillin-resistant *Staphylococcus aureus* (MRSA)



Number per 100 000 population

Reduce by 10% the total incidence of bloodstream infections with third-generation cephalosporin-resistant *Escherichia coli*



Number per 100 000 population

Reduce by 4% the total incidence of bloodstream infections with carbapenem-resistant *Klebsiella pneumoniae*



Number per 100 000 population

2019 baseline	23.65	-
2022	23.58	-0.3%
2030 TARGET	17.3	-27%
2019 baseline	60.4%	-
2022	56.7%	-3.7% *
2030 TARGET	65%	+4.6% *
2019 baseline	4.3	-
2022	3.9	-8.2%
2030 TARGET	3.8	-10%
2019 baseline	7.4	-
2022	7.0	-6.1%
2030 TARGET	6.7	-10%
2019 baseline	1.4	-
2022	3.3	+140%
2030 TARGET	1.3	-4%