



**Reduce by 27% the total consumption of antibiotics in humans**

Defined daily doses (DDDs) per 1 000 inhabitants per day

2019 baseline	30.1	-
2022	33.5	+11.4%
2030 TARGET	22.0	-27%



**At least 65% of the total consumption of antibiotics in humans belongs to the 'Access' group of antibiotics**

As defined in the AWaRe classification of the WHO

2019 baseline	48.9%	-
2022	55.4%	+6.5% *
2030 TARGET	65%	+16.1% *

\*Percentage point difference from 2019.



**Reduce by 18% the total incidence of bloodstream infections with meticillin-resistant *Staphylococcus aureus* (MRSA)**

Number per 100 000 population

2019 baseline	6.9	-
2022	14.6	+113%
2030 TARGET	5.6	-18%

**Reduce by 5% the total incidence of bloodstream infections with third-generation cephalosporin-resistant *Escherichia coli***

Number per 100 000 population

2019 baseline	6.2	-
2022	10.8	+73.6%
2030 TARGET	5.9	-5%

**Reduce by 5% the total incidence of bloodstream infections with carbapenem-resistant *Klebsiella pneumoniae***

Number per 100 000 population

2019 baseline	2.6	-
2022	9.9	+278%
2030 TARGET	2.5	-5%

