



Antimicrobial resistance targets Greece

Target achieved Progress Regress



Reduce by 27% the total consumption of antibiotics in humans

Defined daily doses (DDDs) per 1 000 inhabitants per day

2019
baseline

34.1

-

2022

32.9

-3.5%

2030
TARGET

24.9

-27%

2019
baseline

46.8%

-

2022

42.6%

-4.2%
*

2030
TARGET

65%

+18.2%
*

*Percentage point difference from 2019.



At least 65% of the total consumption of antibiotics in humans belongs to the 'Access' group of antibiotics

As defined in the AWaRe classification of the WHO

2019
baseline

4.6

-

2022

5.0

+8.2%

2030
TARGET

4.1

-10%



Reduce by 10% the total incidence of bloodstream infections with meticillin-resistant *Staphylococcus aureus* (MRSA)

Number per 100 000 population

2019
baseline

2.6

-

2022


4.0

+54.7%

2030
TARGET

2.6

-



Maintain at baseline level the total incidence of bloodstream infections with third-generation cephalosporin-resistant *Escherichia coli*

Number per 100 000 population

2019
baseline

13.1

-

2022

18.0

+38.0%

2030
TARGET

12.4

-5%



Reduce by 5% the total incidence of bloodstream infections with carbapenem-resistant *Klebsiella pneumoniae*

Number per 100 000 population