



# Antimicrobial resistance targets Netherlands

Target achieved Progress Regress



**Reduce by 3% the total consumption of antibiotics in humans**

Defined daily doses (DDDs) per 1 000 inhabitants per day

2019  
baseline

9.5

-

2022

9.1

-4.3%

2030  
TARGET

9.2

-3%



**At least 65% of the total consumption of antibiotics in humans belongs to the 'Access' group of antibiotics**

As defined in the AWaRe classification of the WHO

2019  
baseline

71.2%

-

2022

71%

-0.2%  
\*

2030  
TARGET

65%

-

\*Percentage point difference from 2019.



**Reduce by 3% the total incidence of bloodstream infections with meticillin-resistant *Staphylococcus aureus* (MRSA)**

Number per 100 000 population

2019  
baseline

0.40

-

2022


0.52

+31.7%

2030  
TARGET

0.38

-3%



**Maintain at baseline level the total incidence of bloodstream infections with third-generation cephalosporin-resistant *Escherichia coli***

Number per 100 000 population

2019  
baseline

4.5

-

2022

4.3

-5.4%

2030  
TARGET

4.5

-



**Maintain at baseline level the total incidence of bloodstream infections with carbapenem-resistant *Klebsiella pneumoniae***

Number per 100 000 population

2019  
baseline

0.025

-

2022

0.046

+85.9%

2030  
TARGET

0.025

-