



Reduce by 27% the total consumption of antibiotics in humans

Defined daily doses (DDDs) per 1 000 inhabitants per day

2019 baseline	25.8	-
2022	27.6	+7.0%
2030 TARGET	18.8	-27%



At least 65% of the total consumption of antibiotics in humans belongs to the 'Access' group of antibiotics

As defined in the AWaRe classification of the WHO

2019 baseline	52.8%	-
2022	50.3%	-2.5% *
2030 TARGET	65%	+12.2% *

*Percentage point difference from 2019.



Reduce by 18% the total incidence of bloodstream infections with meticillin-resistant *Staphylococcus aureus* (MRSA)

Number per 100 000 population

2019 baseline	13.7	-
2022	20.3	+48.0%
2030 TARGET	11.3	-18%

Reduce by 5% the total incidence of bloodstream infections with third-generation cephalosporin-resistant *Escherichia coli*

Number per 100 000 population

2019 baseline	6.3	-
2022	10.8	+70.3%
2030 TARGET	6.0	-5%

Reduce by 5% the total incidence of bloodstream infections with carbapenem-resistant *Klebsiella pneumoniae*

Number per 100 000 population

2019 baseline	7.1	-
2022	26.3	+269%
2030 TARGET	6.8	-5%

